

LA-UR-19-22546

Approved for public release; distribution is unlimited.

| | |
|---------------|--|
| Title: | Report Viewgraphs for IC project: Regional ocean and land ice ensemble studies to better quantify uncertainties in Antarctic-driven sea level rise |
| Author(s): | Veneziani, Carmela |
| Intended for: | Report |
| Issued: | 2019-03-20 |

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Report Viewgraphs for IC project: Regional ocean and land ice ensemble studies to better quantify uncertainties in Antarctic-driven sea level rise

- Two sets of simulations were performed within this allocation:
 - 1) a 10-year experiment using the ACCESS 1.3 model data for the open ocean boundary conditions, and MAR-downscaled ACCESS 1.3 data for the atmospheric forcing;
 - 2) a 15-year experiment with similar conditions as the one in 1), but with a different initial condition for temperature and salinity nearby the ice shelves.
- Results from simulation 1) are presented in Figure 1, in terms of sea-ice extent, and they show a significant loss of model sea-ice in winter compared with observations. Further investigation of these model biases revealed an important bias in the MAR-ACCESS precipitation, with the model underestimating precipitation over the ocean to the point that strong convection was triggered. The impact of these biases on the upper ocean stratification appears to eventually affect the sub ice shelf melt rates as well (see Fig. 2).
- Results from simulation 2) are presented in Figure 2 in terms of averaged melt rates for specific ice shelves around Antarctica. The simulated melt rates are comparable with observational values, although a trend is visible and possibly due to the biases described above.




Figure 1: Sea-ice concentration (%) in winter simulated by the ROMS model (left panels; climatologies are computed over years 1984-1989 and for months June-July-August), and observed by SSM/I satellite observations.




Figure 2: Time series of mass loss from ice-shelves (melt rates) in ROMS, for specific ice shelves, and compared with observed values reported in Rignot et al, 2013 (dots and shading).